Mass mortality in Northwestern Mediterranean rocky benthic communities: effects of the 2003 heat wave

J. GARRABOU*'[†], R. COMA[‡], N. BENSOUSSAN*, M. BALLY*, P. CHEVALDONNÉ*, M. CIGLIANO[§], D. DIAZ[¶], J. G. HARMELIN*, M. C. GAMBI[§], D. K. KERSTING[∥], J. B. LEDOUX*, C. LEJEUSNE*, C. LINARES*, C. MARSCHAL*, T. PÉREZ*, M. RIBES[†], J. C. ROMANO^{**}, E. SERRANO[‡], N. TEIXIDO[‡], O. TORRENTS*, M. ZABALA^{††}, F. ZUBERER^{‡‡} and C. CERRANO[§]§

*UMR 6540 – DIMAR CNRS/Université de' Aix-Marseille, Centre d'Océanologie de Marseille, Station Marine d'Endoume, rue Batterie des Lions, 13007 Marseille, France, †Institut de Ciències del Mar CSIC, Passeig Marítim de la Barceloneta 37-49, E-08003 Barcelona, Spain, ‡Centre d'Estudis Avançats de Blanes CEAB-CSIC, Accés Cala St. Francesc 14, 17300 Blanes, Girona, Spain, §Stazione Zoologica 'Anton Dohrn', Villa Comunale, 80121 Napoli, Italy, ¶Instituto Español de Oceanografía, C/Moll de Ponent s/n, 07015 Palma de Mallorca, Spain, ∥Reserva Marina de Las Islas Columbretes, SGPM-Tragsa, Castelló, Spain, **UMR-CNRS 6134, Centre Scientifique de Vignola, route des Sanguinaires, 20000 Ajaccio, France, ††Departament d'Ecologia, Universitat de Barcelona, Avda. Diagonal 645, 08028 Barcelona, Spain, ‡‡UMS 2196 Centre d'Océanologie de Marseille, Station Marine d'Endoume, rue Batterie des Lions, 13007 Marseille, France, §§Dipartimento per lo Studio del Territorio e delle sue Risorse, Università di Genova, C.so Europa 26, 16132 Genova, Italy

Abstract

Late in summer 2003, extensive mass mortality of at least 25 rocky benthic macroinvertebrate species (mainly gorgonians and sponges) was observed in the entire Northwestern (NW) Mediterranean region, affecting several thousand kilometers of coastline. We were able to characterize the mortality event by studying six areas covering the main regions of the NW Mediterranean basin. The degree of impact on each study area was quantified at 49 sites by estimating the proportion of colonies affected in populations of several gorgonian species compared with reference data obtained in years without mortality signs. According to these data, the western areas (Catalan coast and Balearic Islands) were the least affected, while the central areas (Provence coast and Corsica-Sardinia) showed a moderate impact. The northernmost and eastern areas (Gulf of Genoa and Gulf of Naples) displayed the highest impact, with almost 80% of gorgonian colonies affected. The heat wave of 2003 in Europe caused an anomalous warming of seawater, which reached the highest temperatures ever recorded in the studied regions, between 1 and 3 °C above the climatic values (mean and maximum). Because this exceptional warming was observed in the depth ranges most affected by the mortality, it seems likely that the 2003 anomalous temperature played a key role in the observed mortality event. A correlation analysis between temperature conditions and degree of impact seems to support this hypothesis. Under the present climate warming trend, new mass mortality events may occur in the near future, possibly driving a major biodiversity crisis in the Mediterranean Sea.

Keywords: disturbance, effects of climate change, marine biodiversity, mortality outbreak

Received 9 April 2008; revised version received 24 October 2008 and accepted 28 October 2008

Introduction

Recent climatic trends have resulted in significant responses in marine ecosystems, such as pelagic productivity, shifts in species' geographical distributions, community composition changes, and modifications of migratory patterns (Rosenzweig *et al.*, 2007). The increase in mass mortality outbreaks in coastal ecosystems during the past few decades has also been linked to modifications in environmental conditions caused by global change (Harvell *et al.*, 1999; Harley *et al.*, 2006). In coastal ecosystems, coral reefs are suffering a dramatic increase in the frequency and intensity of mortality and

Correspondence: Present address: J. Garrabou, Institut de Ciències del Mar CSIC, Passeig Marítim de la Barceloneta 37-49, E-08003 Barcelona, Spain, tel. + 34 93 230 95 00, fax + 34 93 230 95 55, e-mail: garrabou@icm.csic.es

disease, including several regional-scale mass mortality events in gorgonians (Nagelkerken et al., 1997) and widespread coral bleaching episodes (Glynn, 1991; McWilliams et al., 2005). In general, high-temperature conditions are documented to be concurrent with observations of mass mortalities (Harvell et al., 1999; Lesser et al., 2007). These events have occurred in the context of significant warming of the global average surface (+0.76 °C for the 20th century; IPCC, 2007) and upper global ocean (+0.3 °C from 1950 to 2000; Levitus et al., 2000). Projections for the 21st century depict an increase in warming $(+1.8 \text{ or } +3.5 \degree \text{C} \text{ depending on})$ the emission scenario; IPCC, 2007) along with a very likely increase in the occurrence of heat waves (Déqué 2007; Diffenbaugh et al., 2007). Under these conditions, incidences of mortality events may spread over larger geographic areas and a wider spectrum of organisms. This scenario could cause a marine biodiversity loss crisis. Analyzing the impact of these events at appropriate scales (spatial and temporal) and biological organization levels (species, populations, communities) is crucial to accurately anticipate future changes in marine ecosystems and propose adapted management and conservation plans.

In summer 2003, the European region suffered an exceptional heat wave (Luterbacher *et al.*, 2004; Schär *et al.*, 2004). This had dramatic social, economic, and environmental consequences, including an increase in heat-related human morbidity and mortality, forest fires, and widespread parasites (De Bono *et al.*, 2004). This event began in early June 2003 and lasted until mid-August, with air temperature records about 3-6 °C above the seasonal average in many parts of Europe (Schär *et al.*, 2004). This heat wave also affected the seawater temperature in the Northwestern (NW) Mediterranean region (Marullo & Guarracino, 2003; Sparnocchia *et al.*, 2006). In the marine realm, a large-scale mass mortality event was observed in different areas of the NW Mediterranean during this heat wave.

The NW Mediterranean has been affected by several severe mortality events of benthic invertebrate species during the last few decades (Coma & Ribes, 2003; Garrabou *et al.*, 2003). In 1999, an unprecedented large-scale mass mortality event was recorded, affecting about 30 benthic species over hundreds of kilometers of the NW Mediterranean coast (Cerrano *et al.*, 2000; Perez *et al.*, 2000). Anomalous high-temperature conditions during summer 1999 were identified as the main triggering factor of the mortality (Cerrano *et al.*, 2000; Perez *et al.*, 2000). Bearing in mind the warming trend observed in the Mediterranean (Bethoux *et al.*, 1990; Vargas-Yáñez *et al.*, 2008) and the coincidence of the event with anomalous temperature conditions (Cerrano *et al.*, 2000; Romano *et al.*, 2000), it was hypothesized that new

mass mortality events could occur in the near future (Cerrano *et al.*, 2000; Coma *et al.*, 2000; Perez *et al.*, 2000). Unfortunately, this hypothesis was corroborated in 2003 when a further large-scale mass mortality event was observed in the NW Mediterranean region.

The main goal of this paper is to provide a comprehensive report on the impact of the 2003 mass mortality event in the NW Mediterranean area by providing data on the species affected, the intensity of the impact, the depth range affected, and the timing of the event for different NW Mediterranean areas. The affected communities have been studied during the last four decades by the staff of various research institutes. Therefore, even though we cannot provide quantitative data for all species, we contend that our observations on unusual mortality signs allow us to provide an unbiased assessment of the impact of the 2003 mass mortality event. Finally, data on temperature conditions during the summer of 2003 are also reported to analyze the potential role of anomalous temperature conditions in the occurrence of the event.

Materials and methods

Study areas

This study encompasses a large geographic scale, covering the main rocky coasts over several thousands of kilometers of coastline from 39°50′N 0°40′E in Spain to 40°34′N 14°20′E in Italy, including the main Mediterranean islands (Balearic, Corsica, and Sardinia; Fig. 1). In an attempt to provide data on the extent and impact of the 2003 mass mortality event at the regional scale, the NW Mediterranean region has been divided into six main study areas, each spanning hundreds of kilometers of coastline (Fig. 1). A total of 49 sites were surveyed throughout the study. For each area, between 4 and 14 sites were surveyed (Table 1).

For each area, we collated reports of the 2003 mass mortality events for the rocky coasts collected by various research teams from Spain, France, and Italy. These included reports of park officers of different marine protected areas and diving clubs operating in the study areas.

Impact assessment of the 2003 mortality event

Data on mortality impact were based on qualitative and quantitative surveys. In the qualitative surveys, lists of species showing clear-cut signs of recent necrosis, such as denuded skeletons in gorgonians, sponges, and scleractinian corals and empty valves attached to the substratum in bivalves, were acquired for each study site. All species included in the qualitative surveys were large, macrobenthic, easily identifiable species, and



Fig. 1 Northwestern Mediterranean showing the limits of the six study areas where surveys of the 2003 mass mortality were conducted.

often abundant at the surveyed sites. The depth range affected by mortality and data on the timing of the event were provided for each site.

The quantitative surveys focused on gorgonian species because (a) they were abundant at all study areas and are considered engineering species in Mediterranean communities, (b) the effects of mortality are easy to quantify, and (c) gorgonian populations have been monitored at several study sites for up to 15 years, so they provide an excellent baseline that is not available for other affected species (Harmelin & Marinopoulos, 1994; Garrabou & Harmelin, 2002; Coma *et al.*, 2004; Linares *et al.*, 2008). Four gorgonian species were considered in assessing the quantitative impact of the 2003 mass mortality event: the red gorgonian *Paramuricea clavata*, the yellow gorgonian *Eunicella cavolinii*, the white gorgonian *Eunicella singularis*, and the red coral *Corallium rubrum*.

Surveys quantified the percentage of recent tissue necrosis (i.e., denuded axis or recent epibiosis) in about 100 colonies for each species present at each study site and depth range. We considered a colony to be affected by the mortality when it showed recent tissue necrosis over $\geq 10\%$ of its surface. For each survey, the percentage of affected colonies was calculated as an indicator of the mortality impact. This indicator was chosen instead of the mean extent of necrosis or the whole mortality rate for two main reasons. First, the mean extent of tissue necrosis could introduce a potential bias in a large-scale assessment, because it integrates the effects of past disturbances suffered by gorgonian populations (e.g., Harmelin & Marinopoulos, 1994; Coma et al., 2004; see 'Discussion'). Second, previous studies showed that the immediate impact of the mass mortality events is much smaller in terms of total colony mortality than partial mortality (Cerrano et al., 2005; Linares et al., 2005).

Surveys were carried out at the end of 2003 (September to November) just after the observation of the mortality event. In 2003, a total of 93 surveys (more than 10 000 colonies observed) were carried out in the six study areas (Table 2a). A Kruskal–Wallis analysis was used to test for the differences in the impact among study areas (including all gorgonians species considered in each study area). Multiple comparisons were also performed to determine specific interarea differences. We used nonparametric analysis because the data followed neither normality nor homogeneity of variances after different transformations.

We ultimately compared the results of the 2003 surveys with surveys carried out in 2002, 2004, and 2005 (unaffected years), for which a total of 36 surveys (more than 6000 colonies observed) were performed on the Catalan coast, Provence coasts, and Corsica-Sardinia study areas (Table 2b). In this comparison, we only used surveys carried out on the red gorgonian *P. clavata*. For this species, the dataset was the largest available, allowing a reliable evaluation of the impact of the 2003 mortality event. Data from unaffected years was pooled and compared with the 2003 data using a Kolgomorov-Smirnov test. As previously, we used nonparametric analysis because the data followed neither normality nor homogeneity of variances after different transformations. The Kolgomorov-Smirnov test and the Kruskal-Wallis test were computed using the program STATISTICA (version 8.0, StatSoft, Inc., Tulsa, OK, USA).

Temperature conditions during summer 2003

Temperature records from 1, 10, and 20 m depth were analyzed to determine the temperatures that occurred during the mass mortality of invertebrates at the end of summer 2003. A total of six temperature series were

	Study areas								
Impact characteristics	1 – Balearic Islands	2 – Catalan coast	3 – Provence coast	4 – Gulf of Genoa	5 – Corsica and Sardinia	6 – Gulf of Naples			
Number of surveyed sites per area	8	9	14	4	10	4			
Maximum depth affected (m)	30-40	15-20	15–25	25-30	25–30	25-30			
Total number of affected species	5	3	16	14	13	5			
Affected species									
Gorgonians									
Paramuricea clavata	А	А	А	А	А	А			
Eunicella cavolinii	NO	NO	А	А	А	А			
Eunicella sigularis	А	А	А	А	NO	А			
Eunicella verrucosa	NO	NO	NA	А	NO	NO			
Leptogorgia sarmentosa	NO	NO	NA	А	NO	NO			
Corallium rubrum	NA	А	А	А	А	NA			
Scleractinians									
Cladocora caespitosa	А	NA	NA	А	NO	А			
Balanophyllia europaea	NO	NO	NO	А	NO	А			
Zoantharians									
Parazoanthus axinellae	NA	NA	NA	А	NA	NA			
Sponges									
Spongia officinalis	NO	NA	А	А	А	NA			
Ircinia dendroides	NO	NA	А	NO	NO	NO			
Ircinia variabilis	А	NA	А	А	А	NO			
Ircinia oros	NO	NA	NA	А	NA	NO			
Agelas oroides	NO	NA	А	А	NA	NA			
Crambe crambe	NA	NA	А	NA	А	NA			
Cacospongia spp.	NO	NA	А	NO	А	NO			
Hippospongia communis	NO	NA	А	NO	А	NO			
Petrosia ficiformis	NO	NA	А	А	А	NA			
Bryozoans									
Myriapora truncata	А	NA	NA	NA	NA	NA			
Sertella spp.	NO	NA	NA	NA	А	NA			
Bivalves									
Spondylus gaederopus	NA	NA	А	А	А	NA			
Lima spp.	NA	NA	NO	NO	А	NA			
Arca spp.	NA	NA	А	А	А	NA			
Ostrea edulis	NO	NA	А	NO	NO	NA			
Lithophaga lithophaga	NA	NA	А	NO	NO	NO			

Table 1	Impact characteristics and	species affected	during the 2003 mass	s mortality event in the six s	tudy areas

For each affected species, we provide information regarding the impact for each study area: A, affected; NA, not affected; NO, no observations available.

analyzed, some from national observatories and some collected by the research teams. These series provided temperature data for all of the study areas except for the Corsica-Sardinia area, for which temperature records were not available. The main features of the temperature series are shown in Table 3. The series of L'Estartit and Villefranche cover the last 30 years and are the longest temperature records for the NW Mediterranean. Not all depths were available for each site: only temperatures at 1 m were available at Columbretes (Balearic Island area), while the 10 m level is missing at L'Estartit (Catalan coast area). The six temperature records were collected using

various measurement strategies, although in general, temperature measurements were derived from routine measurements (daily, weekly, and biweekly) carried out near the coast. Average mean and maximum temperatures from 1 July to 30 September were calculated for each available year from the data series to examine the temperature anomaly of 2003 within each temperature record.

Temperature and mortality

A correlation analysis between the 2003 summer temperature conditions (TXS_{2003}) and the percentage of

1094 J. GARRABOU *et al.*

Table 2 (a) Mean, minimum, and maximum percentages of colonies with $\geq 10\%$ of colony surface showing recent necrosis (denuded axis or recent epibiosis) obtained from (1) *Paramuricea clavata*, (2) *Eunicella singularis*, (3) *Eunicella cavolinii*, and (4) *Corallium rubrum* gorgonian populations dwelling in different areas of the Northwestern (NW) Mediterranean. Surveys were carried out in late 2003 in the depth range affected by the 2003 mass mortality event. (b) Mean percentage of colonies with $\geq 10\%$ of colony surface showing recent necrosis (denuded axis or recent epibiosis) obtained from *Paramuricea clavata* populations dwelling in different areas of the Northwestern (NW) Mediterranean in years not affected by mass mortality events

					% affected colon	ies
Study area	N surveys	s N coloni	es Dep	oth range	$X \pm SD$	Minimum–maximum
(a)						
(1) Paramuricea clavata						
Balearic Islands*	3	316	30-	40	14.9 ± 23.4	0-41.8
Catalan coast	15	1067	15–2	20	4.8 ± 6.3	0-21.2
Provence coast	15	1680	5—2	25	28.5 ± 21.5	1.9-77.7
Gulf of Genoa	4	300	25-3	30	82.5 ± 9.5	70–90
Corsica-Sardinia	6	1417	15–2	25	54.4 ± 22.1	19.6-84.9
Gulf of Naples	3	165	25-	32	64.9 ± 15.0	12.5-100
(2) Eunicella singularis						
Balearic Islands	5	896	15–1	20	9.5 ± 4.5	3.1-14.3
Catalan coast	4	349	10-	15	2.7 ± 2.4	0–5.7
Provence coast	-	_	-		-	-
Gulf of Genoa	2	100	25-	30	85 ± 7.07	80-90
Corsica-Sardinia	_	_	-		-	-
Gulf of Naples	6	329	15–2	22	86.6 ± 12.3	11.1-100
(3) Eunicella cavolinii						
Balearic Islands	-	_	-		-	-
Catalan coast	_	_	_		_	-
Provence coast	11	1147	15–1	25	14.5 ± 14.5	3.0-50.8
Gulf of Genoa	1	100	10-	30	90	-
Corsica-Sardinia	7	848	15–1	20	17.1 ± 10.6	4.9-34.2
Gulf of Naples	_	_	_		-	-
(4) Corallium rubrum						
Balearic Islands	-	_	-		-	-
Catalan coast	_	_	-		-	-
Provence coast	8	1140	5–2	25	33.5 ± 28.0	5.7-91.4
Gulf of Genoa	1	300	25-	30	50	-
Corsica-Sardinia	2	399	20-2	25	20.2 ± 17.9	7.6-32.9
Gulf of Naples	_	-	_		-	_
					% affected	colonies
Study area	N surveys	Survey years	N colonies	Depth range	$\overline{X \pm SD}$	Minimum–maximum
(b)						
Paramuricea clavata						
NE Spain coast	16	2002	1644	15–35	0	-
Provence coast	12	2004-2005	3142	15–35	1.9 ± 1.6	0-5.2
Corsica	8	2005	1577	18–25	0.5 ± 0.8	0–2.0
All areas	36		6363		0.8 ± 1.3	0–5.2

*In the Balearic Islands area, *P. clavata* was only affected in several sites from the Columbretes Marine Reserve (situated midway between Balearic Island and the continental coast of the Iberian Peninsula).

affected colonies (see above) explored the potential role of temperature in the mass mortality of invertebrates. The analysis was performed using only data from the Provence area, because in this area the 2003 surveys cover affected and unaffected depth ranges (45 surveys observing more than 5000 colonies) and two temperature data series (Marseilles and Villefranche) were available (Table 3). We applied the nonparametric cor-

Site	Study area	Period	Measurement depths (m)	Sampling frequency	Instrumentation	References
Columbretes	Balearic Islands	1991–2006	1	Daily	Mercury-in-glass thermometer	-
L'Estartit	Catalan coast	1975–2004	1, 20, 50, 80	Weekly	Reversing thermometers	Pascual <i>et al.</i> (1995)
Marseilles	Provence coast	1999–2004	1, 10, 15, 20, 25, 30, 35, 40, 45, 50	Biweekly– monthly	1994–2003: CTD ME ECO-57 2003–2004: SBE 911	Romano <i>et al.</i> (2000)
Villefranche	Provence coast	1975–2004	1, 5, 10, 15, 20, 25, 30, 35, 40, 50, 75	Weekly	1975–1995: Reversing thermometers 1995–2004: CTD SBE 911	Prieur (2002)
Genoa	Gulf of Genoa	1994–2004	1, 10, 20, 50, 80	Biweekly– monthly	ODAS buoy and Idronaut Ocean Seven probe	Siccardi <i>et al.</i> (1998), Schiaparelli <i>et al.</i> (2007)
Naples	Gulf of Naples	2001–2004	1, 10, 20, 30 up to 80	Weekly	CTD-Seabird 911 plus	Ribera d'Alcalà <i>et al</i> . (2004)

Table 3 Main features of temperature data series analyzed to determine the 2003 positive temperature anomaly in the Northwestern Mediterranean basin

relation analysis (Spearman rank order, STATISTICA version 8.0, StatSoft, Inc.) because the percentage of affected colonies followed neither normality nor homogeneity of variances after different transformations.

Results

Temperature conditions during summer 2003

The six temperature series analyzed indicated positive temperature anomalies at 1 and 10 m depth in all study areas. The analysis of the longest data series available from the Catalan coast (L'Estartit, Spain, 1975-present), and Provence (Villefranche, France, 1975-present) showed that the 2003 summer was the warmest of the last 30 years. In general, at 1 and 10 m depth, the mean and maximum temperatures during the 2003 summer were greater than the mean values (Table 4). In the 2003 summer, temperature (TXS₂₀₀₃) varied between 23.8 and 27.5 °C, yielding differences between 0.7 and 2.8 °C in relation to average values (TXS). Regarding temperature maxima, Tmax₂₀₀₃ varied between 25.1 and 29.6 °C, representing 1.2 and 2.9 °C differences compared with the average maximum temperatures (Tmax) (Table 4). The Gulf of Genoa showed the highest deviations from the mean values (Table 4).

At 20 m depth, the 2003 mean summer temperatures (TXS_{2003}) were similar or lower than the mean values (TXS). However, in most areas where mortality was observed beyond 20 m, temperature maxima $(Tmax_{2003})$ were always more than 1 °C greater than the average maximum temperatures (Tmax), except in the Gulf of

Naples where the 2003 temperature maximum was about 0.5 °C lower than the average value (Table 4).

Qualitative surveys: species and depth range affected and timing of the mortality

Overall, 25 benthic species were affected in the NW Mediterranean region by the 2003 mass mortality event (Table 1). Most species showed signs of unusual total and/or partial mortality over large geographic areas (several hundreds of kilometers; Table 1 and Fig. 2). The affected species belonged to four phyla: Cnidaria, Porifera, Bryozoa, and Mollusca. All species are hard substrate dwellers and several are prominent components of the infralittoral and circalittoral communities (photophilic algae assemblages, precoralligenous, coralligenous, and semidark caves). The total number of affected species per study area varied from a minimum of three to five species in the southern areas to a maximum of 12-16 species in the northern ones. The depth range affected was limited to 15-30 m, except in the Balearic Islands area where effects could be detected up to 35-40 m. From the Catalan coast to the Gulf of Naples, the depth range affected increased west to east (Table 1).

In most areas, the first signs of mortality were observed at the end of August 2003. At that time, for the gorgonian *P. clavata*, colony parts showing necrosis displayed a change in color from red to gray, resulting from the decomposition of the organic parts of the cortex, leaving only sclerite aggregations. The remaining organic parts and sclerites then progressively detached from the colony's axis, resulting in the partial and total denudation of skeletons (Fig. 2a). As in pre-

Site	Study area	Period	TXS	Tmax	TXS ₂₀₀₃	Tmax ₂₀₀₃
1 m						
Columbretes	1	1991-2006	25.9 ± 0.8	27.5 ± 0.9	26.7	29.6
L'Estartit	2	1975-2004	21.9 ± 0.7	24.0 ± 1.1	23.2	25.4
Marseilles	3	1999-2004	21.3 ± 1.5	24.4 ± 1.2	23.1	26.5
Villefranche	3	1975-2004	23.6 ± 0.8	24.6 ± 1.0	25.3	26.5
Genoa	4	1994-2004	23.6 ± 0.8	25.5 ± 1.3	26.4	27.8
Naples	6	2001-2004	25.6 ± 1.5	27.3 ± 0.9	26.3	28.6
10 m						
Marseilles	3	1999-2004	20.7 ± 1.7	23.9 ± 1.2	22.4	25.1
Villefranche	3	1975-2004	22.7 ± 0.8	23.8 ± 0.9	24.3	26.2
Genoa	4	1994-2004	22.8 ± 0.9	24.7 ± 1.8	25.2	27.6
Naples	6	2001-2004	24.6 ± 1.7	26.5 ± 1.3	25.3	28.3
20 m						
L'Estartit	2	1975-2004	20.2 ± 0.6	22.5 ± 1.0	20.8	23.3
Marseilles	3	1999-2004	19.5 ± 1.8	22.9 ± 1.9	19.9	21.8
Villefranche	3	1975-2004	21.1 ± 0.9	22.5 ± 0.9	21.3	23.8
Genoa	4	1994-2004	22.1 ± 1.7	22.4 ± 1.6	20.6	23.5
Naples	6	2001-2004	21.1 ± 2.3	23.9 ± 0.9	21.2	23.3

Table 4 Temperature records (from 1 July to 30 September) at 1, 10, and 20 m depths obtained from each study area except for theCorsica-Sardinia area for the years indicated under 'Period'

Two series (Marseilles and Villefranche) were analyzed for the Provence coast. Records were available at 1, 10, and 20 m depth at all sites, except Columbretes (only 1 m) (Balearic Islands study area) and L'Estartit (only 1 and 20 m) (Catalan coast area). TXS and Tmax refer to the average mean and maximum temperatures, respectively, obtained for the available period in each study area. TXS₂₀₀₃ and Tmax₂₀₀₃ refer to the average and maximum temperatures obtained for summer 2003.

vious events, this process could be sudden, with populations shifting from healthy to severely damaged over a few days. Once colonies lost their affected tissue, bare axial skeleton portions became colonized in the following weeks by macroscopic pioneer taxa such as hydroids, bryozoans, and serpulid polychaetes. At the end of September and early October, most gorgonian colonies showed some degree of recent epibiosis on their denuded axis, and thereafter, no further signs of mortality were observed, at least in these species. A similar pattern of tissue loss was found in the other affected gorgonian species (Fig. 2b and c). The scleractinian coral Cladocora caespitosa suffered extensive tissue loss directly through tissue necrosis or after suffering bleaching, resulting in the total or partial deaths of colonies (Fig. 2f). In sponges, specimens also showed total and/ or partial death (Fig. 2d and e). The main affected sponge species were Spongia officinalis, Petrosia ficiformis, Cacospongia spp., and Ircinia spp. Skeletons in horny sponge species remained attached to the substratum as proof of recent mortality (Fig. 2e). Healed wounds were observed in late autumn 2003 in some specimens affected by partial mortality. In bivalve mollusk species, the presence of empty valves attached to the substratum indicated the abnormal recent deaths of individuals. High levels of recent injuries were also noted in some bryozoan species (Myriapora truncata and Sertella spp.).

Quantitative surveys

In 2003, gorgonian populations displayed a dramatic impact with an increase of about 5–82 fold of affected colonies compared with years without mortality for which values around 1% were found (Table 2b; Kolgomorov–Smirnov test, P < 0.001).

The degree of impact varied from about 5% in the least impacted area (Catalan coast) to 75-80% in the most impacted (Gulf of Genoa and Gulf of Naples) (Fig. 3). The Balearic Islands, Provence coast, and Corsica-Sardinia showed intermediate impact values of 15-35% (Fig. 3). These differences were statistically significant (Kruskal–Wallis test, H = 57.39, df = 5, P < 0.001), and overall, the multiple comparisons test supported the three degrees of impact observed. Interspecific differences in the degree of impact were also clear (Fig. 3). The red gorgonian, P. clavata, was among the most affected species in all areas, reaching up to an average of 80% of colonies affected in the Gulf of Genoa (Fig. 3 and Table 2a). The white and yellow gorgonian species, E. singularis and E. cavolinii, were also severely affected on the Italian coasts (Gulf of Genoa and Gulf of Naples), with values up to 80% (Fig. 3 and Table 2a). Meanwhile, in the other areas, both Eunicella species showed a low degree of impact (Fig. 3). For instance, only about 2% of *E. singularis* colonies were affected on the Catalan coast.



Fig. 2 Species affected by the 2003 mass mortality event. (a) *Paramuricea clavata* colonies showing almost complete loss of tissue (denuded axis, light brown areas), with only the tips of branches still alive (red parts) (19 m depth in the Réserve Naturelle de Scandola, Corsica-Sardinia). (b) *Eunicella cavolinii* showing recent signs of partial mortality (pale parts of the axis) at 25 m depth in the Portofino Promontory (Gulf of Genoa). (c) A red coral *Corallium rubrum* colony with signs of partial mortality (dark red areas) contrasting with live tissue (pale red areas) (Réserve Naturelle de Scandola, Corsica-Sardinia). (d) *Petrosia ficiformis* at 8 m depth showing partial mortality (white parts) (Gulf of Genoa). (e) *Spongia officinalis* displaying partial mortality (brown parts) (Réserve Naturelle de Scandola, Corsica-Sardinia). (f) *Cladocora caespitosa*: detail of polyps suffering tissue necrosis (Columbretes Islands Marine Reserve, Balearic Islands).

Finally, the red coral, *C. rubrum*, showed a variable response depending on the study area (Fig. 3 and Table 2a).

According to the number of affected species and the degree of impact in gorgonian populations, the lowest impact was found in the western areas, while the eastern ones showed the highest impact (Table 1 and Fig. 3).

Temperature and mortality

The Spearman rank order correlation analysis between the TXS₂₀₀₃ and the percentage of affected colonies showed a significant association (Fig. 4; r = 0.51, P < 0.05). The higher the temperature was, the higher was the impact on the populations (Fig. 4). The lowest values corresponded to surveys at depths unaffected by the mortality.



Fig. 3 Mean values of the percentage of colonies affected by the 2003 mass mortality event found in each study area considering data for all species surveyed and for each species (*Paramuricea clavata, Eunicella singularis, Eunicella cavolinii,* and *Corallium rubrum*).



Fig. 4 Relationship between 2003 summer temperature conditions TXS₂₀₀₃ (mean 1 July to 30 September) and percentage of affected colonies in the Provence area observed in late 2003. Temperature data from Marseilles and Villefranche from 10, 15, 20, 25, and 30 m depth. Each data point represents data from one survey and the TXS₂₀₀₃ corresponding to the sampling depth.

Discussion

Bearing in mind the number of affected species, the degree of impact, and the geographic extent, the 2003 mass mortality can be considered the largest mortality event ever to have occurred in the NW Mediterranean during the last few decades. Except for the 1999 event, all previous events affected a small number of species generally in a small spatial extent (hundreds of meters to a few kilometers of coastline). In 1999, a diversity of species similar to that of 2003 (30 species in 1999; Perez

et al., 2000) were affected but over a smaller geographic area. Besides the areas affected in 1999 (Cerrano et al., 2000; Perez et al., 2000; Coma et al., 2006), the 2003 event also affected the Catalan coast, Corsica-Sardinia, and the Gulf of Naples. Both events occurred at the end of the summer (Cerrano et al., 2000; Perez et al., 2000). Finally, the 2003 mass mortality event was restricted to shallower waters than in the previous 1999 event, when the maximum depth reached 40 m in most parts of the affected areas (Table 1; Cerrano et al., 2000; Perez et al., 2000; Coma et al., 2006). The geographic extent and the number of species affected by the 1999 and 2003 events in the NW Mediterranean region are comparable with some regional-scale bleaching events documented in the Caribbean and other tropical areas (McWilliams et al., 2005).

Most of the species affected by the 2003 mortality event already suffered damages in previous events. At least four of the affected gorgonian species (P. clavata, E. singularis, E. cavolinii, and C. rubrum) had already suffered extensive damage in some areas of the Mediterranean during the last 20 years (Cerrano et al., 2000, 2005; Perez et al., 2000; Garrabou et al., 2001; Linares et al., 2005; Coma et al., 2006; Gambi et al., 2006; Sbrescia et al., 2008). Among other anthozoan affected species, the scleractinian C. caespitosa had previously suffered severe bleaching during 1997–1999 (Perez et al., 2000; Rodolfo-Metalpa et al., 2000). Likewise, commercial sponges (Spongia spp., Hippospongia communis, Ircinia spp.) had been previously affected by a wide mortality outbreak in 1986 and 1999 as well as in other minor events (Vacelet, 1994; Cerrano et al., 2000; Perez et al., 2000). Among bryozoan species, M. truncata was affected during the 1999 event (Perez et al., 2000). Meanwhile, the number of bivalve species affected increased dramatically from two to six during the 1999 and 2003 events, respectively. Lima spp. was the only species that displayed signs of mortality during these two major events. Furthermore, in the 1980s, Lima spp. and other species (Spondylus gaederopus and oysters) suffered extensive mortality in the Provence area (Meinesz & Mercier, 1983). The three Tunicata (solitary ascidian species) Microcosmus spp., Halocynthia papillosa and Pyura dura (Perez et al., 2000), and the coralline red algae Lithophyllum stictaeforme (Cerrano et al., 2000), which suffered extensive damage in 1999, showed no signs of mortality in 2003. In summary, anthozoans, sponges, and bivalves were among the most affected groups in the 2003 mass mortality event as in other mortality events observed in the Mediterranean during the last decades.

In this study, we provided for the very first time a comprehensive report on the impact of a mortality event that occurred in the NW Mediterranean. Comparison with previous studies is not straightforward because different methods were used to assess the impact of events. Bearing in mind that current changes in the ecosystems concern large scales, we urgently need to develop standardized protocols able to efficiently evaluate the effects of disturbances over large geographic extents. Here, the use of the percentage of affected gorgonian colonies provided an excellent descriptor to quantify the impact of the mass mortality event at the regional scale. Evaluation of the status of populations through the use of this kind of indicator should be performed annually to obtain appropriate baseline data and expanded to groups other than gorgonians when possible. Only through the deployment of this strategy can we accurately evaluate the impacts of disturbances associated with global climate change and provide crucial data to analyze the potential factors associated with these dramatic events.

Potential causes of the 2003 mortality event

The analysis of surface sea temperature (SST) (data from 1985 to 2003) confirmed exceptional warming throughout the whole NW Mediterranean region (Marullo & Guarracino, 2003). However, the temperature anomaly was limited to shallow layers (<20 m depth) due to the calm weather conditions between June and August (Sparnocchia *et al.*, 2006). This could explain the differences in the temperature anomaly observed between 20 m and the other two shallower depths (Table 4).

In general, the depth range where signs of mortality were observed corresponded to the depths where extreme temperatures were recorded (Tables 1 and 4). Because the signs of mortality occurred just after the unusually warm period, the 2003 anomalous temperature likely played a key role in this widespread mortality event. The result of the correlation analysis seems to support this hypothesis (Fig. 4). In contrast to 2003, temperatures recorded during summer 1999 were not exceptional. However, in late summer 1999, the water column showed a very high stability with values around 24 °C, resulting in an anomalous warming of 2-3 °C in the water layer overlying the thermocline accompanied by its deepening (up to 30-40 m) (Romano et al., 2000). This exceptional stratification may explain the differential depth range affected in both mortality events.

Exposure to anomalous high temperatures may produce several effects. Exposure to lethal temperatures acts directly on organisms causing metabolic dysfunctions (Torrents *et al.*, 2008). Exposure to sublethal temperatures can also act directly causing physiological stress that may result in an energy shortage due to increased respiration rate (Coma *et al.*, 2002) and/or decrease the efficacy of defense (Cossins & Bowler, 1987; Coma & Ribes, 2003) and indirectly trigger the development of pathogens (Toren *et al.*, 1998; Banin *et al.*, 2001, 2003). In fact, the bacteria *Vibrio corallilyticus* has been recently identified as a causative thermodependent pathogen from diseased *P. clavata* colonies sampled in 2003 (Bally & Garrabou, 2007). However, microorganisms found on colonies during the 1999 gorgonian mortality event appeared to be opportunistic rather than disease-causing (Cerrano *et al.*, 2000; Martin *et al.*, 2002).

Besides anomalous temperature conditions, several putative agents have been identified or evoked as the cause of mortality outbreaks in the Mediterranean, including pathogens for sponges and sea urchins (Vacelet *et al.*, 1994; Boudouresque & Verlaque, 2001), drops in salinity for gorgonian species (Bavestrello *et al.*, 1994), and the development of filamentous algae (Mistri & Ceccherelli, 1996; Schiaparelli *et al.*, 2007). However, high temperatures were a common environmental condition in most events. Finally, global warming has been associated with the increase in different mortality outbreaks in marine species such as seagrasses, fishes, corals, and sea urchins worldwide (Harvell *et al.*, 1999; Atrill *et al.*, 2004; Lesser *et al.*, 2007).

Differential mortality at the intraspecific and interspecific levels, across spatial scales from meters to hundreds of kilometers, was a common phenomenon for the 2003 mass mortality event (Figs 3 and 4). Similar observations were made for previous events in the Mediterranean (Cerrano et al., 2000; Perez et al., 2000) and worldwide, such as extensive bleaching of coral species in different parts of the world (Hughes et al., 2003; McWilliams et al., 2005). These observations provide clear and empirical evidence for the absence of a single mass mortality threshold for species and populations, which may be modulated by both physical and biological factors. For instance, in the Gulf of Naples, differential mortality could be attributed to the high variability in the local hydrographic conditions (Ribera d'Alcalà et al., 2004), as proven in the case of bleaching events (Nakamura & van Woesik, 2001), while among biological factors, adaptation mechanisms could play a key role in the response of populations to temperature stress (Torrents et al., 2008). Despite the complexity underlying the biological response, two major trends can be anticipated regarding the potential impact of warming in the NW Mediterranean region. First, bearing in mind the hydrodynamic conditions in the region, shallow populations (0-50 m) will be most exposed to future mortality events. Second, populations of affected species dwelling in northern areas may be more vulnerable than southern ones, because they are probably adapted to colder temperature conditions and are found in shallower waters

(Zabala & Ballesteros, 1989; Linares *et al.*, 2008). Overall, research on the environmental, physiological, or genetic determinants of this differential response should be considered a major challenge for the coming years. Revealing these factors will help determine how species will cope with future environmental changes and how humans can undertake actions to promote the resilience of affected communities (Hughes *et al.*, 2003).

Consequences of mass mortality events

Most of the affected species are characterized by high longevity and slow population dynamics with low growth, recruitment, and mortality rates (Pansini & Pronzato, 1990; Garrabou & Harmelin, 2002; Coma et al., 2004; Linares et al., 2008). Thus, a dramatic increase in mortality rates (total and partial), as well as short times between mortality events, may produce severe consequences for the viability of most affected populations (Linares et al., 2007). Data on the impact and recovery in gorgonian and sponge populations from the 1999 mass mortality events showed a dramatic decrease in the biomass of populations and a very slow recovery during the 4 years following the impact (Cerrano et al., 2005; Linares et al., 2005; Coma et al., 2006; J. Garrabou et al., unpublished data). Similar patterns were found for populations surveyed after the 2003 event (Cigliano & Gambi, 2007; J. Garrabou et al., unpublished data), except in populations from the Catalan coast, which showed only a low impact. In this area, recovery was complete in the year following the event (R. Coma, unpublished data).

Because several species have suffered damage across a large geographic extent, concerns regarding their recruitment or spill-off from other areas, and therefore their persistence, are justified and large-scale conservation plans are urgently needed. Finally, it must be taken into account that (1) species considered in this study are large, prominent, and easily identifiable, but that small, cryptic, or poorly known species are also affected by such large-scale events (Chevaldonné & Lejeusne, 2003; Scinto *et al.*, 2007); and (2) species such as large sponges, erect bryozoans, and gorgonians provide biogenic structure, so other species not directly affected by the mass mortality but structurally dependent on affected larger taxa could also indirectly suffer the impact through modification of habitat conditions (e.g., flow regime, food availability, shelter) (Soulé et al., 2003; Ribes & Coma, 2005).

Summary and predictions

Projections available for the Mediterranean basin region for the 21st century indicate warmer and drier conditions with a dramatic increase in the frequency and persistence of extreme events such as heat waves (Stott et al., 2004; Déqué 2007; Diffenbaugh et al., 2007; IPCC, 2007). For instance, the number of heat wave days may be multiplied by 10 by the end of the current century (Déqué 2007). Although the relationship between weather and oceanographic conditions is not straightforward, long-term studies on seawater temperature conditions already show a clear-cut warming trend of coastal waters worldwide (Southward et al., 1995, 2005; Levitus et al., 2000; Vargas-Yáñez et al., 2008; J.C. Romano et al., unpublished data). In addition, a recent study on oceanographic conditions of the Mediterranean Sea during the 21st century supports a clear-cut picture of warming, with a nearly spatially homogenous increase in SST of +3.1 °C in 2099 (Somot et al., 2006). These results agree reasonably well with the warming trends calculated from coastal time series over the past three decades (Vargas-Yáñez et al., 2008; J.C. Romano et al., unpublished data).

In this context, the repetition of large-scale mortality events can be expected during the next decades. Consequences of these events could severely expose populations of affected species to local ecological extinction processes, threatening the conservation of the rich Mediterranean biodiversity (Bianchi & Morri, 2000; Ballesteros, 2006; Bianchi, 2007). This climate change is the latest form of human disturbance affecting the Mediterranean coastal communities at the regional scale through mortality outbreaks and shifts in species distributions (Francour et al., 1994; Bianchi & Morri, 2000). The expansion and intensification of climate change effects together with the effects of other strong disturbances (pollution, mechanical habitat destruction, invasive species, and overfishing) may act together, driving the Mediterranean to a major biodiversity crisis during the 21st century (Bianchi & Morri, 2000; Jackson et al., 2001; Sala, 2004).

Acknowledgements

The authors gratefully acknowledge the assistance during sampling of J.M. Dominici (Réserve Naturelle de Scandola, Parc Regional de Corse), and B. de Ligondes and R. Graille (Centre d'Océanologie de Marseille). Data on seawater temperatures were provided by J. Pasqual (Catalan coast), Service d'Observation en Milieu Littoral, INSU-CNRS, Observatoire Océanologique de Villefranche sur mer and Centre d'Océanologie de Marseille (Provence coast) and by Drs Vincenzo Saggiomo and Maurizio Ribera D'Alcalà from the Stazione Zoologica 'Anton Dohrn' of Naples (Gulf of Naples). This research was partly supported by the IFB Foundation (Institut Français de la Biodiversité), the environmental department of TOTAL-Fina-Elf group, an 'Action Concertée Iniciative' (ACI) program 'Jeunes chercheurs 2003' (number JC5360) from the French Research Ministry, research grants CTM2006-01463 and CGL2007-66757 from the 'Ministerio de Educación y Ciencia' of Spain and the Medchange project funded by the Agence Nationale pour la Recherche (ANR).

References

- Atrill MJ, Kelmo F, Jones MB (2004) Impact of the 1997–98 El Niño event on the coral reef-associated echinoderm assemblage from northern Bahia, northeastern Brazil. *Climate Research*, 26, 151–158.
- Ballesteros E (2006) Mediterranean coralligenous assemblages: a synthesis of present knowledge. *Oceanography and Marine Biology: An Annual Review*, **44**, 123–195.
- Bally M, Garrabou J (2007) Thermodependent bacterial pathogens and mass mortalities in temperate benthic communities: a new case of emerging disease linked to climate change. *Global Change Biology*, **13**, 2078–2088.
- Banin E, Israely T, Fine M, Loya Y, Rosenberg E (2001) Role of endosymbiotic zooxanthellae and coral mucus in the adhesion of the coral-bleaching pathogen *Vibrio shiloi* to its host. *FEMS Microbiology Letters*, **199**, 33–37.
- Banin E, Vassilakos D, Orr E, Martinez RJ, Rosenberg E (2003) Superoxide dismutase is a virulence factor produced by the coral bleaching pathogen *Vibrio shiloi*. *Current Microbiology*, 46, 418–422.
- Bavestrello G, Bertone S, Cattaneo-Vietti R, Cerrano C, Gaino E, Zanzi D (1994) Mass mortality of *Paramuricea clavata* (Anthozoa: Cnidaria) on Portofino Promontory cliffs (Ligurian Sea). *Marine Life*, 4, 15–19.
- Bethoux JP, Gentili B, Raunet J, Tailliez D (1990) Warming trend in the western Mediterranean deep water. *Nature*, **347**, 660–662.
- Bianchi CN (2007) Biodiversity issues for the forthcoming tropical Mediterranean Sea. *Hydrobiologia*, 580, 7–21.
- Bianchi CN, Morri C (2000) Marine biodiversity of the Mediterranean Sea: situation, problems and prospects for future research. *Marine Pollution Bulletin*, 40, 367–376.
- Boudouresque CF, Verlaque M (2001) Ecology of Paracentrotus lividus. In: Ecology of Paracentrotus lividus. Edible Sea Urchins: Biology and Ecology (ed. Lawrence JM), pp. 177–215. Elsevier, London, UK.
- Cerrano C, Arillo A, Azzini F *et al.* (2005) Gorgonian population recovery after a mass mortality event. *Aquatic Conservation*, **15**, 147–157.
- Cerrano C, Bavestrello G, Bianchi CN *et al.* (2000) A catastrophic mass-mortality episode of gorgonians and other organisms in the Ligurian Sea (northwestern Mediterranean), summer 1999. *Ecology Letters*, **3**, 284–293.
- Chevaldonné P, Lejeusne C (2003) Regional warming-induced species shift in north-west Mediterranean marine caves. *Ecology Letters*, **6**, 371–379.
- Cigliano M, Gambi MC (2007) The long hot summer. A further mortality event of gorgonians along the Phlaegrean Islands (Tyrrhenian Sea). *Biologia Marina Mediterranea*, **14**, 292–293.
- Coma R, Linares C, Ribes M, Diaz D, Garrabou J, Ballesteros E (2006) Consequences of a mass mortality in populations of *Eunicella singularis* (Cnidaria: Octocorallia) in Menorca (NW Mediterranean). *Marine Ecology Progress Series*, **327**, 51–60.

- Coma R, Pola E, Ribes M, Zabala M (2004) Long-term assessment of temperate octocoral mortality patterns, protected vs. unprotected areas. *Ecological Applications*, 14, 1466–1478.
- Coma R, Ribes M (2003) Seasonal energetic constraints in Mediterranean benthic suspension feeders: effects at different levels of ecological organization. *Oikos*, **101**, 205–215.
- Coma R, Ribes M, Gili JM, Zabala M (2000) Seasonality in coastal benthic ecosystems. Trends in Ecology and Evolution, 15, 448–453.
- Coma R, Ribes M, Gili JM, Zabala M (2002) Seasonality of in situ respiration rate in three temperate benthic suspension feeders. *Limnology and Oceanography*, **47**, 324–331.
- Cossins AR, Bowler K (1987) *Temperature Biology of Animals*. Chapman & Hall, London, UK.
- De Bono A, Pedruzzi P, Giuliani G, Kluser S (2004) *Impacts of summer 2003 heat wave in Europe*. Early Warning on Emerging Environmental Threats, UNEP/DEWA/ ~ Europe/GRID-Geneva, 4 pp.
- Déqué M (2007) Frequency of precipitation and temperature extremes over France in an anthropogenic scenario: model results and statistical correction according to observed values. *Global and Planetary Change*, **57**, 16–26.
- Diffenbaugh NS, Pal JS, Giorgi F, Gao XJ (2007) Heat stress intensification in the Mediterranean climate change hotspot. *Geophysical Research Letters*, **34**, L11706, 1–6.
- Francour P, Boudouresque CF, Harmelin JG, Harmelin-Vivien ML, Quignard JP (1994) Are the Mediterranean waters becoming warmer? Information from biological indicators. *Marine Pollution Bulletin*, 28, 523–526.
- Gambi MC, Cigliano M, Iacono B (2006) Segnalazione di un evento di mortalità di gorgonacei lungo le coste delle isole di Ischia e Procida (Golfo di Napoli, Mar Tirreno). *Biologia Marina Mediterranea*, **13**, 583–587.
- Garrabou J, Harmelin JG (2002) A 20-year study on life-history traits of a harvested long-lived temperate coral in the NW Mediterranean: insights into conservation and management needs. *Journal of Animal Ecology*, **71**, 966–978.
- Garrabou J, Perez T, Chevaldonné P *et al.* (2003) Is global change a real threat for conservation of the NW Mediterranean marine biodiversity? *Geophysical Research Abstracts*, **5**, 10522.
- Garrabou J, Perez T, Sartoretto S, Harmelin JG (2001) Mass mortality event in red coral *Corallium rubrum* populations in Provence region (France, NW Mediterranean). *Marine Ecology Progress Series*, 217, 263–272.
- Glynn PW (1991) Coral reef bleaching in the 1980s and possible connections with global warming. *Trends in Ecology and Evolution*, **6**, 175–179.
- Harley CDG, Hughes AR, Hultgren KM *et al.* (2006) The impacts of climate change in coastal marine systems. *Ecology Letters*, **9**, 228–241.
- Harmelin JG, Marinopoulos J (1994) Population structure and partial mortality of the gorgonian *Paramuricea clavata* in the north-western Mediterranean. *Marine Life*, **4**, 5–13.
- Harvell CD, Kim K, Burkholder JM et al. (1999) Emerging marine diseases – climate links and anthropogenic factors. Science, 285, 1505–1510.
- Hughes TP, Baird AH, Bellwood DR *et al.* (2003) Climate change, human impacts, and the resilience of coral reefs. *Science*, **301**, 929–933.

© 2009 The Authors

Journal compilation © 2009 Blackwell Publishing Ltd, Global Change Biology, 15, 1090–1103

1102 J. GARRABOU *et al.*

- IPCC (2007) Climate change 2007: the physical science basis. Contribution of working group I to the Fourth Assessment Report of the intergovernmental panel on climate change (eds Solomon SD, Qin M, Manning Z *et al.*,) Cambridge University Press, Cambridge, UK and New York, NY, USA, 996p.
- Jackson JBC, Kirby MX, Berger WH *et al.* (2001) Historical overfishing and the recent collapse of coastal ecosystems. *Science*, 293, 629–638.
- Lesser MP, Bythell JC, Gates RD, Johnstone RW, Hoegh-Guldberg O (2007) Are infectious diseases really killing corals? Alternative interpretations of the experimental and ecological data. *Journal of Experimental Marine Biology and Ecology*, **346**, 36–44.
- Levitus S, Antonov JI, Boyer TP, Stephens C (2000) Warming of the World Ocean. *Science*, **5461**, 2225–2229.
- Linares C, Coma R, Diaz D, Zabala M, Hereu B, Dantart L (2005) Immediate and delayed effects of a mass mortality event on gorgonian population dynamics and benthic community structure in the NW Mediterranean Sea. *Marine Ecology Pro*gress Series, **305**, 127–137.
- Linares C, Coma R, Garrabou J, Diaz D, Zabala M (2008) Size distribution, density and disturbance of two Mediterranean gorgonians: *Paramuricea clavata* and *Eunicella singularis*. *Journal* of Applied Ecology, 45, 688–699.
- Linares C, Doak DF, Coma R, Diaz D, Zabala M (2007) Life history and viability of a long-lived marine invertebrate: the octocoral *Paramuricea clavata*. *Ecology*, **88**, 918–928.
- Luterbacher J, Dietrich D, Xoplaki E, Grosjean M, Wanner H (2004) European seasonal and annual temperature variability, trends and extremes since 1500. *Science*, **303**, 1499–1503.
- Martin Y, Bonnefort JL, Chancerelle L (2002) Gorgonians mass mortality during the 1999 late summer in French Mediterranean coastal waters: the bacterial hypothesis. *Water Research*, **36**, 779–782.
- Marullo S, Guarracino M (2003) L'anomalia termica del 2003 nel Mar Mediterraneao ossevata da satellite. *Energia, Ambiente e Innovazione*, **6**, 48–53.
- McWilliams JP, Côté IM, Gill JA, Sutherland WJ, Watkinson AR (2005) Accelerating impacts of temperature-induced coral bleaching in the Caribbean. *Ecology*, **86**, 2055–2060.
- Meinesz A, Mercier D (1983) Sur les fortes mortalities de Spondylus gaederopus Linné observées sur les côtes de Méditerranée. Travaux Scientifiques du Parc National de Port-Cros, 9, 89–95.
- Mistri M, Ceccherelli VU (1996) Effects of a mucilage event on the Mediterranean gorgonian *Paramuricea clavata*. Short term impacts at the population and colony levels. *Italian Journal of Zoology*, **63**, 221–230.
- Nagelkerken I, Buchan K, Smith GW *et al.* (1997) Widespread disease in Caribbean sea fans: II. Patterns of infection and tissue loss. *Marine Ecology Progress Series*, **160**, 255–263.
- Nakamura T, van Woesik R (2001) Water-flow rates and passive diffusion partially explain differential survival of corals during the 1998 bleaching event. *Marine Ecology Progress Series*, **212**, 301–304.
- Pansini M, Pronzato R (1990) Observations on the dynamics of a Mediterranean sponge community. In: *New Perspectives in*

Sponge Biology (ed. Rützler K), pp. 404–415. Smithsonian Institution Press, Washington, DC, USA.

- Pascual J, Salat J, Palau M (1995) Evolución de la temperatura del mar entre 1974 y 1994, cerca de la costa catalana. *Internationale Colloque Okeanos*, **95**, 23–28.
- Perez T, Garrabou J, Sartoretto S, Harmelin JG, Francour P, Vacelet J (2000) Mass mortality of marine invertebrates: an unprecedented event in the Northwestern Mediterranean. *Comptes Rendus de l'Académie des Science Paris, Science de la Vie,* **323**, 853–865.
- Prieur L (2002) Physical historical data on the Ligurian Sea from the Villefranche-sur-mer Observatory. In: *Tracking Long-Term Hydrological Change in the Mediterranean Sea*. CIESM Workshop Series, no. 16, pp. 37–40.
- Ribera d'Alcalà M, Conversano F, Corato F *et al.* (2004) Seasonal pattern in plankton communities in a pluriannual time series at the coastal Mediterranean site (Gulf of Naples): an attempt to discern recurrences and trends. *Scientia Marina*, **68**, 65–83.
- Ribes M, Coma R (2005) The role of engineer species in the benthic–pelagic coupling: the study case of a Mediterranean gorgonian. *Limnology and Oceanography*, Summer meeting, 128 (Abstract).
- Rodolfo-Metalpa R, Bianchi CN, Peirano A, Morri C (2000) Coral mortality in NW Mediterranean. *Coral Reefs*, **19**, 24.
- Romano JC, Bensoussan N, Younes WAN, Arlhac D (2000) Anomalies thermiques dans les eaux du golfe de Marseille durant l'été 1999. Une explication partielle de la mortalité d'invertébrés fixés. Comptes Rendus de l'Académie des Science Paris, Science de la Vie, 323, 415–427.
- Rosenzweig C, Casassa G, Karoly DJ et al. (2007) Assessment of observed changes and responses in natural and managed systems. Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change (eds Parry ML, Canziani OF, Palutikof JP, van der Linden PJ, Hanson CE), pp. 79–131. Cambridge University Press, Cambridge, UK.
- Sala E (2004) The past and present topology and structure of Mediterranean subtidal rocky-shore food webs. *Ecosystems*, 7, 333–340.
- Sbrescia L, Russo M, Di Stefano F, Russo GF (2008) Analisi delle popolazioni di gorgonie dell'AMP di Punta Campanella dopo eventi di moria di massa. Proceedings Associazione Italiana di Oceanologia e Limnologia. Atti Associazione Italiana di Oceanologia e Limnologia, **19**, 449–457.
- Schär C, Vidale PL, Lüthi D, Frei C, Häberli C, Liniger M, Appenzeller C (2004) The role of increasing temperature variability in European summer heat waves. *Nature*, **427**, 332–336.
- Schiaparelli S, Castellano M, Povero P, Sartoni G, Cattaneo-Vietti R (2007) A benthic mucilage event in North-Western Mediterranean Sea and its possible relationships with the summer 2003 European heatwave: short term effects on littoral rocky assemblages. *Marine Ecology an Evolutionary Perspective*, **28**, 341–353.
- Scinto A, Benvenuto C, Cerrano C, Mori M (2007) Seasonal cycle of Jassa marmorata Holmes, 1903 (Crustacea, Amphipoda) in

the Ligurian Sea (Mediterranean Sea, Italy). *Journal of Crustacean Biology*, **27**, 212–216.

- Siccardi A, Orsi M, Centurioni L (1998) La boa ODAS Italia 1, un centro di misura integrato in Mar Ligure. In: Atti del 12° Congresso dell'Associazione Italiana di Oceanologia e Limnologia, Vol. 2 (ed. Piccazzo M), AIOL, Genova, Italy pp. 263–272.
- Somot S, Sevault F, Déqué M (2006) Transient climate change scenario simulation of the Mediterranean Sea for the twenty-first century using a high-resolution ocean circulation model. *Climate Dynamics*, **27**, 851–879.
- Soulé ME, Estes JA, Berger J, Del Rio CM (2003) Ecological effectiveness: conservation goals for interactive species. *Con*servation Biology, 17, 1238–1250.
- Southward AJ, Hawkins SJ, Burrows MT (1995) 70 years observations of changes in distribution and abundance of zooplankton and intertidal organisms in the Western English Channel in relation to rising sea temperature. *Journal of Thermal Biology*, 20, 127–155.
- Southward AJ, Langmead O, Hardman-Mountford NJ *et al.* (2005) Long-term oceanographic and ecological research in the Western English Channel. *Advances in Marine Biology*, **47**, 1–105.
- Sparnocchia S, Schiano ME, Picco P, Bozzano R, Cappelletti A (2006) The anomalous warming of summer 2003 in the surface layer of the Central Ligurian Sea (Western Mediterranean). *Annual Geophysics*, **24**, 443–452.

- Stott PA, Stone DA, Allen R (2004) Human contribution to the European heatwave of 2003. *Nature*, **432**, 610–614.
- Toren A, Landau L, Kushmaro A, Loya Y, Rosenberg E (1998) Effect of temperature on adhesion of *Vibrio* strain AK-1 to *Oculina patagonica* and on coral bleaching. *Applied Environmental Microbiology*, **64**, 1379–1384.
- Torrents O, Tambutte E, Caminiti N, Garrabou J (2008) Upper thermal thresholds of shallow *vs.* deep populations of the precious Mediterranean red coral *Corallium rubrum* (L.): assessing the potential effects of warming in the NW Mediterranean. *Journal of Experimental Marine Biology and Ecology*, **357**, 7–19.
- Vacelet J (1994) The struggle against the epidemic which is decimating Mediterranean sponges. FAO Technical Report, FAO, Rome.
- Vacelet J, Gaino E, Gallissian M, Vacelet E (1994) Bacterial attack of spongin skeleton during the 1990 Mediterranean sponge disease. In: Sponges in Time and Space; Biology, Chemistry, Paleontology (eds van Soest RWM, van Kempen TMG, Braekman JC), pp. 355–362. Balkema, Rotterdam, the Netherlands.
- Vargas-Yáñez M, Garcia MJ, Salat J, Garcia-Martinez MC, Pascual J, Moya F (2008) Warming trends and decadal variability in the Western Mediterranean shelf. *Global and Planetary Change*, 63, 177–184.
- Zabala M, Ballesteros E (1989) Surface-dependent strategies and energy flux in benthic marine communities or, why corals do not exist in the Mediterranean. *Scientia Marina*, **53**, 3–17.