

AN ANNOTATED LIST OF ALIEN MARINE SPECIES OF THE ISCHIA ISLAND (GULF OF NAPLES)

This contribution was presented as a poster at the BioMarine Workshop “Management of Bioinvasions in the Mediterranean Sea - a way forward”, held on the island of Ischia on 4-5 May 2016, with the aim of updating the information and providing the workshop’s participants with a list of the alien taxa that have been recorded in the area hosting the meeting.

The island of Ischia, and the nearby islands of Procida and Vivara (Phlaegean islands), are located on the northern side of the Gulf of Naples, and are included within the Marine Protected Area of the “Regno di Nettuno” (“Neptune’s Realm”) since April 2008 (Gambi *et al.*, 2003).

The Phlaegean islands are considered a biogeographic boundary which constitutes the distribution limit of most South Mediterranean thermophilous species, since the 14 °C *minimum winter surface temperature* isotherm (the 14 °C divide) is occurring between Ischia and the Pontine archipelago (e.g., Ventotene island) (Gambi *et al.*, 2009; Gambi, 2014). In the past two decades, due to global warming also detected in the Mediterranean surface waters, we observed a northward shift of the 14 °C divide, while the minimum winter isotherm in the area of the Phlaegean islands is now often rising to the 15 °C divide. Therefore this zone is an interesting observatory for the distribution and migratory pattern of thermophilous species both native and introduced/aliens (Gambi, 2014). Occurrence and increase in the abundance of some native Mediterranean warm-species have been documented for this area (Dappiano and Gambi, 2004; Cigliano *et al.*, 2007; Gambi, 2014), as for the coral *Astroides calycularis* (Pallas, 1776), the bryozoan *Calpensia nobilis* (Esper, 1796), and the fishes *Sparisoma cretense* (L., 1758), *Sphoeroides pachygaster* (Müller & Troschel, 1848) and *Sphyraena viridensis* Cuvier, 1829.

As for the alien taxa the annotated list of the species recorded at Ischia is reported in Tab. 1.

A total of 22 alien species have been documented including 1 dinoflagellate, 4 macroalgae, 1 sponge, 3 hydrozoans, 7 polychaetes, 2 molluscs, 2 crustaceans, 1 bryozoan and 1 fish. Although some of the listed taxa are occasional and have been recorded in the area only once (e.g., the fish *F. commersonii* or the opistobranch *B. leachi*), this number of aliens is relatively high, in comparison with areas of larger extension. As an example, 47 taxa are reported from the Central Tyrrhenian Sea and 31 from the Southern Tyrrhenian Sea (see Occhipinti Ambrogi *et al.*, 2011). The high number of records from Ischia is probably in relation to the large amount of benthic studies issued from the activity of the Stazione Zoologica, especially at the Villa Dohrn-Benthic ecology Center (Ischia), whose investigations around the island date back to the late 1960s (Gambi, 2014).

Some of the species recorded show an invasive habit, such as the dinoflagellate *O. ovata*, the macroalgae *C. cylindracea* and *A. armata*, and the invertebrates *B. bairdi*, *P. paucibranchiata*. Most of the species have a Indo-Pacific origin and some of them likely entered from the Red Sea via the Suez Canal (e.g., *B. luctuosum*, *B. leachi*, *F. commersonii*), while *P. magna*, *B. bairdi* and *A. verticillata* have a South Atlantic or Caribbean origin.

Finally, it is interesting to note that various alien species have been recorded from the acidified zones (low and extreme low pH conditions) of the Castello Aragonese CO₂ vents, such as *O. ovata*, *A. taxiformis*, *C. cylindracea*, *B. bairdi*, *N. infratorquata*, *Mesanthura* sp., *P. gibbesi* (Fig. 1), and thus in agreement with what observed with other “nuisance” species in other CO₂ vent’s systems (Hall-Spencer and Allen, 2015). Several other species were reported inside or in the vicinity of the harbor of Ischia, e.g., *S. comatus*, *B. bairdi*, *P. antennata*, or associated to artificial structures, e.g., *P. magna*, *B. luctuosum*, *B. bairdi* and *A. verticillata*, thus confirming the general observation that more impacted and low competition habitats are more prone to the colonization/invasion by aliens (Occhipinti Ambrogi *et al.*, 2011).

Tab. 1

Alien species/origin	Notes and references
(*) <i>Ostreopsis ovata</i> Fukuyo, 1981 (Dinoflagellata, Dinophyceae) (Indo-Pacific)	Recorded in Ischia since 2011-2012 (reported in Di Cioccio <i>et al.</i> , 2014), including the acidified areas of the Castello CO ₂ vents. Recorded also recently in September 2016 (Gambi M.C., pers. observ.)
(*) <i>Caulerpa cylindracea</i> (Sonder) Verlaque, Huisman & Boudouresque (Chlorophyceae) (Indo-Pacific)	Recorded since 2000 in various areas off Ischia (Buia <i>et al.</i> , 2003 indicated as <i>Caulerpa racemosa</i>) is highly invasive especially in summer and autumn. It occurs also on the CO ₂ vents off the Castello Aragonese (Hall-Spencer <i>et al.</i> , 2008) although not all years (Gambi M.C., pers. observ.)
(*) <i>Asparagopsis taxiformis</i> (Delile) Trevisan de Saint-Leon (Rhodophyceae) (Indo-Pacific)	Recorded in 2000 off Ischia and Procida (Flagella <i>et al.</i> , 2003; D'Archino <i>et al.</i> , 2003 in Gambi <i>et al.</i> , 2003) and confirmed to belong to <i>A. taxiformis</i> based on genetic analysis by Andreakis <i>et al.</i> (2004). At present, it is very abundant/invasive in spring also in the north acidified area of the Castello Aragonese CO ₂ vents, both on bare rock and among <i>Posidonia</i> shoots (Gambi M.C., person. observ.)
<i>Acrothamnion preissii</i> (Sonder) E.M. Wollaston (Rhodophyceae) (Indo-Pacific)	Recorded in June 2015 in shallow rocky reefs (10 m depth) off Ischia (Ballesteros E., Teixido N. & Gambi M.C., unpublished data)
<i>Womersleyella setacea</i> (Hollenberg) R.E. Norris (Rhodophyceae) (Indo-Pacific)	Recorded in June 2015 in shallow rocky reefs (10 m depth) off Ischia (Ballesteros E., Teixido N. & Gambi M.C., unpublished data)
<i>Paraleucilla magna</i> Klatau <i>et al.</i> , 2004 (Porifera) (South Atlantic, Brazil)	Recorded in 2014 on artificial barriers and structures around the coast of Ischia (Grech <i>et al.</i> , 2015)
<i>Eudendrium merulum</i> Watson, 1985 (Hydrozoa) (South Pacific)	Observed along the coast of Ischia by Piraino S. (pers. observ.) and reported in Gravili <i>et al.</i> (2010)
<i>Clytia linearis</i> (Thorneley, 1900) (Hydrozoa) (Circumtropical)	Recorded as epiphytic on <i>Posidonia oceanica</i> by Boero (1981a, b)
<i>Cirrholaenia tetraneura</i> Kramp, 1959 (Hydrozoa) (Indo-Pacific)	Recorded by Brinckmann (1965), but already signaled at Ischia by Uchida (1964) and identified as <i>Eugymnanthea minuta</i> (see Kubota, 1995)
(*) <i>Pseudopolydora paucibranchiata</i> Okuda, 1937 (Polychaeta, Spionidae) (Indo-Pacific)	Recorded in July 2014 in the soft bottoms of the harbor of Ischia, but present in the harbor since 1977 (identified as <i>Polydora antennata</i> in Fresi <i>et al.</i> , 1983, 1984). The present record represents the fist in the Western Mediterranean (Radashevsky V., Giangrande A. & Gambi M.C., unpublished data)
<i>Lysidice collaris</i> Grube, 1870 (Polychaeta, Eunicidae) (Red Sea, lessepsian)	Species borer of <i>Posidonia oceanica</i> seagrass sheaths, recorded in Ischia since 1995 (Guidetti <i>et al.</i> , 1997)
<i>Streblosoma comatus</i> (Grube, 1859) (Polychaeta, Terebellidae) (Indo-Pacific)	Reported as <i>Streblosoma hesslei</i> in the harbor of Ischia in 1977 (Giangrande <i>et al.</i> , 1981)
<i>Branchiomma luctuosum</i> (Grube, 1869) (Polychaeta, Sabellidae) (Red Sea, lessepsian)	Recorded in Ischia in 1992 on a <i>Cymodocea nodosa</i> meadow (Sordino and Gambi, 1992) and in artificial moorings (Gambi M.C., pers. observ.); it seems no more present in this area (Gambi M.C., pers. observ.).
(*) <i>Branchiomma bairdi</i> (McIntosh, 1885) (Polychaeta, Sabellidae) (Caribbean Sea)	Recorded in July 2012 inside the marinas of Lacco Ameno, Casamicciola and Ischia. It occurs also in the acidified areas of the south side CO ₂ vents of the Castello (Arias <i>et al.</i> , 2013), where is now becoming invasive (Gambi M.C., pers. observ.)
<i>Novafabricia infratorquata</i> (Fitzhugh, 1973) (Polychaeta, Fabriciidae) (Caribbean Sea)	Recorded in rocky vegetated reef of the low pH area of the Castello CO ₂ vent's system in 2008 (Giangrande <i>et al.</i> , 2014)

<i>Spirorbis marioni</i> Caulery & Mesnil, 1897 (Polychaeta, Serpulidae) (Indo-Pacific)	Recorded in 2008 on artificial settlement structures off the Castello Aragonese area at Ischia but in areas with normal pH conditions (Cigliano <i>et al.</i> , 2010)
<i>Percnon gibbesi</i> (H. Milne Edwards, 1853) (Crustacea Decapoda) (Atlantic and Pacific)	Visually observed since 2004 in several intertidal/upper subtidal areas of Ischia (Lorenti M., Zupo V., Gambi M.C., pers. observ.; Dappiano and Gambi, 2004). A few juveniles, identified as <i>P. cf. gibbesi</i> , have been collected in the <i>Posidonia oceanica</i> meadow off the Castello d'Ischia in 2011 (Garrard <i>et al.</i> , 2014)
<i>Mesanthura</i> sp. (Crustacea, Isopoda) (Indo-Pacific?)	A species showing the traits of an alien, likely conspecific with <i>M. romulea</i> Poore & Lew Ton, 1986 (Indo-Pacific) (Lorenti <i>et al.</i> , 2009). Found on the rocky reefs off the Castello Aragonese at Ischia (Kroeker <i>et al.</i> , 2011)
<i>Bursatella leachi</i> De Blainville, 1817 (Mollusca Opisthobranchia) (Red Sea, lessepsian)	Recorded at Ischia in 1986 (Russo, 1987) in a <i>Cymodocea nodosa</i> meadow. Never recorded after
<i>Polycerella emertoni</i> Verrill, 1881 (Mollusca Opistobranchia) (Amphi-Atlantic species)	Recorded as <i>Polycerella recondita</i> Schmekel, 1965 by Schmekel (1968) in the Cartaromana Bay at Ischia (2-10 m depth). Never reported after
(*) <i>Amathia (Zoobotryon) verticillata</i> (Delle Chiaje, 1822) (Bryozoa) (Caribbean Sea)	Observed in 2000 as epiphytic on <i>Cymodocea nodosa</i> and some macroalgae and gorgonians (Gambi M.C., pers. observ.), and in more recent years (2013-2014) in artificial reefs off Ischia (Grech <i>et al.</i> , 2015)
<i>Fistularia commersonii</i> Rueppell, 1838 (Osteichthyes) (Red Sea, lessepsian)	Recorded at Ischia (Sant'Angelo cliff, 14 m depth) in November 2014 (Tiberti <i>et al.</i> , 2015)

(*) = invasive

Acknowledgements – We wish to thank Piraino S. and Gravili C. for providing information on the Hydrozoa, and Procaccini G., Andreakis N., Crocetta F., and Giangrande A. for insights on some of the species. Giuseppe Renella filmed the video of the fish *F. commersonii* at S. Angelo (Ischia).

References

- ANDREAKIS N., PROCACCINI G., KOOISTRA W.H.C.F. (2004) - *Asparagopsis taxiformis* and *Asparagopsis armata* (Bonnemaisoniales, Rhodophyta): genetic and morphological identification of Mediterranean populations. *Europ. Phycol. J.*, **39** (3): 273-283.
- ARIAS A., GIANGRANDE A., GAMBI M.C., ANADON N. (2013) - Biology and new records of the invasive species *Branchiomma bairdi* (Annelida: Sabellidae) in the Mediterranean Sea. *Mediterr. Mar. Sci.*, **14** (1): 162-171.
- BOERO F. (1981a) - Bathimetric distribution of the epifauna of a *Posidonia* meadow of the Isle of Ischia (Naples): Hydroids. *Rapp. P.-v. Réun. Commn Int. Explor. Scient. Mer. Méditerr.*, **27** (2): 197-198.
- BOERO F. (1981b) - Systematics and ecology of the hydroid population of two *Posidonia oceanica* meadows. *Mar. Ecol.*, **2** (3): 181-197.
- BRINCKMANN A. (1965) - The life cycle of the medusa *Cirrhoholovenia tetraneura* Kramp, 1959 (Leptomedusae, Lovenellidae) with a hydroid of the genus *Cuspidella* Hincks. *Can. J. Zool.*, **43**: 13-15.
- BUIA M.C., GAMBI M.C., DAPPIANO M. (2003) - La distribuzione dell'alga alloctona *Caulerpa racemosa* (Forsskål) J. Agardh (Chlorophiceae) nelle isole Flegree. *Acc. Sc. Lett. Arti Napoli, Mem. Soc. Sc. Fis. Mat.*, **5**: 201-212.
- CIGLIANO M., COCITO S., GAMBI M.C. (2007) - Epibiosis of *Calpensia nobilis* (Esper) (Bryozoa: Cheilostomida) on *Posidonia oceanica* (L.) Delile rhizomes: effects on borer colonization and morpho-chronological features of the plant. *Aquat. Bot.*, **86** (1): 30-36.
- CIGLIANO M., GAMBI M.C., RODOLFO-METALPA R., PATTI F.P., HALL-SPENCER J.M. (2010) - Effects of ocean acidification on invertebrate settlement at natural volcanic CO₂ vents. *Mar. Biol.*, **157** (11): 2489-2502.
- DAPPIANO M., GAMBI M.C. (2004) - New data on occurrence of thermophile scleractinia (Cnidaria, Anthozoa) in the Phlaegean islands (Ischia, Procida, Vivara - Gulf of Naples), with special attention to *Astroides calicularis*. In: Marine Biogeography of the Mediterranean Sea: selected topics. *Biogeographia*, **25**: 31-46.

DI CIOCCIO D., BUIA M.C., ZINGONE A. (2014) - Ocean acidification will not deliver us from *Ostreopsis*. In: Kim H.G., Reguera B., Hallegraeff G.M., Lee C.K., Han M.S., Choi J.K. (eds), *Harmful Algae 2012. Proceedings of the 15th International Conference on Harmful Algae*: 85-88.

FLAGELLA M.M., GUALA I., ESPOSITO A., LORENTI M., BUIA M.C. (2003) - *Asparagopsis taxiformis* and *Caulerpa racemosa*: ecophysiological traits and interaction with native algal communities In: Proceedings of the 3rd European Phycological Congress, Belfast (UK), 21-26 July, Book of Abstract: 37 pp.

FRESI E., COLOGNOLA R., GAMBI M.C., GIANGRANDE A., SCARDI M. (1983) - Ricerche sui popolamenti bentonici di substrato duro del Porto d'Ischia. Infralitorale fotofilo: Policheti I. *Cah. Biol. Mar.*, **24**: 1-19.

FRESI E., COLOGNOLA R., GAMBI M.C., GIANGRANDE A., SCARDI M. (1984) - Ricerche sui popolamenti di substrato duro del Porto d'Ischia. Infralitorale fotofilo. Policheti II. *Cah. Biol. Mar.*, **25** (1): 33-47.

HALL-SPENCER J.M., ALLEN R. (2015) - The impact of CO₂ emissions on "nuisance" marine species. *Res. Reports Biodiv. Studies*, **4**: 33-46.

HALL-SPENCER J.M., RODOLFO-METALPA R., MARTIN S., RANSOME E., FINE M., TURNER S.M., ROWLEY S.J., TEDESCO D., BUIA M.C. (2008) - Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature*, **454**: 96-99.

KROEKER K.J., MICHELI F., GAMBI M.C., MARTZ T.R. (2011) - Divergent ecosystem responses within a benthic marine community to ocean acidification. *Proc. Natl. Acad. Scie USA*, **108** (35): 14515-14520.

KUBOTA S. (1995) - Cnidome and growth of a medusa of *Cirrholovenia tetranema* (Leptomedusae, Cirrholeniidae) in Japan. *Publ. Seto Mar. Bioi. Lab.*, **36** (5-6): 365-378.

GAMBI M.C. (2014) - L'isola d'Ischia: un osservatorio speciale per lo studio del cambiamento climatico globale a mare. In: Leone U., Greco P. (eds), *Ischia Patrimonio dell'Umanità. Natura e cultura*. Doppia voce Ed., Napoli: 71-97.

GAMBI M.C., BARBIERI F., BIANCHI C.N. (2009) - New record of the alien seagrass *Halophila stipulacea* (Hydrocharitaceae) in the Western Mediterranean: a further clue to changing Mediterranean Sea biogeography. *Mar. Biodiv. Rec.*, **2** (e84): 1-7. doi:10.1017/S175526720900058X

GAMBI M.C., DE LAURO M., JANNUZZI F. (eds) (2003) - Ambiente marino costiero e territorio delle isole Flegree (Ischia Procida Vivara - Golfo di Napoli). Risultati di uno studio multidisciplinare. *Acc. Sc. Lett. Arti Napoli, Mem. Soc. Sc. Fis. Mat.*, **5**: 425 pp.

GARRARD S., GAMBI M.C., SCIPIONE M.B., PATTI F.P., LORENTI M., ZUPO V., PATERSON D.M., BUIA M.C. (2014) - Indirect effects may buffer negative responses of seagrass invertebrate communities to ocean acidification. *J. Exp. Mar. Biol. Ecol.*, **461**: 31-38.

GIANGRANDE A., GAMBI M.C., FRESI E. (1981) - Two species of Polychaetes new to the Mediterranean fauna. *Boll. Zool.*, **48** (2-3): 311-317.

GIANGRANDE A., GAMBI M.C., MICHELI F., KROEKER K.J. (2014) - Fabriciidae (Annelida, Sabellida) from a naturally acidified coastal system (Italy) with description of two new species. *J. Mar. Biol. Assoc. UK*, **94** (7): 1417-1427.

GRAVILI C., BELMONTE G., CECERE E., DENITTO F., GIANGRANDE A., GUIDETTI P., LONGO C., MASTROTOTARO F., MOSCATELLO F., PETROCELLI A., PIRAINO S., TERLIZZI A., BOERO F. (2010) - Nonindigenous species along the Apulian coasts, Italy. *Chem. Ecol.*, **26** (Suppl.): 121-142.

GRECH D., PATTI F.P., CHIARORE A., MULAS M., BUIA M.C. (2015) - Coastal transformation and marine habitat loss. In: Ozhan E. (ed), *Proc. 12th MEADCOAST*, Varna Bulgaria: 271-281.

GUIDETTI P., BUSSOTTI S., GAMBI M.C., LORENTI M. (1997) - Invertebrate borers in *Posidonia oceanica* scales: relationships between their distribution and lepidochronological parameters. *Aquat. Bot.*, **58** (2): 151-164.

LORENTI M., DAPPIANO M., GAMBI M.C. (2009) - Occurrence and ecology of *Mesanthura* (Crustacea: Isopoda: Anthuridea) in two Italian harbours. *Mar. Biodiv. Rec.*, **2** (e48). DOI: 10.1017/S1755267209000542

OCCHIPINTI AMBROGI A., MARCHINI A., CANTONE G., CASTELLI A., CHIMENZ C., CORMACI M., FROGLIA C., FURNARI G., GAMBI M.C., GIACCONE G., GIANGRANDE A., GRAVILI C., MASTROTOTARO F., MAZZIOTTI C., ORSI RELINI L., PIRAINO S. (2011) - Alien species along the Italian coasts: an overview. *Biol. Invasions*, **13**: 215-237. doi:10.1007/s10530-010-9803-y

RUSSOG.F.(1987) - Segnalazione di *Bursatella leachi* De Blainville, 1817 (Mollusca, Opisthobranchia, Aplysiomorpha) per le acque dell'isola d'Ischia e considerazioni sull'ecologia della specie. *Boll. Soc. Naturalisti in Napoli*, **94**: 243-253.

SCHMEKEL L. (1968) - Ascoglossa, Notaspidea und Nudibranchia im Litoral des Golfes von Neapel. *Rev. Suisse Zool.*, **65**: 103-155.

SORDINO P., GAMBI M.C. (1992) - Prime osservazioni sulla biologia riproduttiva e sul ciclo vitale di *Branchiomma luctuosum* (Grube, 1869) (Polychaeta, Sabellidae). *Oebalia*, (Suppl.) **17**: 425-427.

TIBERTI L., BARBIERI F., GAMBI M.C. (2015) - Osservazioni sulla presenza di specie aliene lungo le coste della Campania: un esempio di conoscenza ecologica locale e “cittadinanza scientifica”. *Notiziario SIBM*, **67** (2015): 23-29 (on line su www.sibm.it).

UCHIDA T. (1964) - Medusae of *Eugymnanthea*, an epizoic hydroid. *Publ. Seto Mar. Bio. Lab.*, **12** (1): 101-107.

Maria Cristina GAMBI

Maurizio LORENTI

Francesco Paolo PATTI

Valerio ZUPO

*Stazione Zoologica ‘Anton Dohrn’
Dept. Integrative Marine Ecology
Villa Dohrn, Benthic Ecology Center
Ischia (NA), Italy.*



Fig. 1 - Some of the alien species recorded along the coast of Ischia: top left) *Caulerpa cylindracea* settled among *Posidonia* at the Castello south side low pH zone (3 m depth; photo: Iacono B.); top right) *Asparagopsis taxiformis* settled on bare rocks at the Castello north side low pH zone (3 m depth; photo: Gambi M.C.); bottom left) *Brachiomma bairdi* settled at the Castello south side low pH zone (1 m depth; photo: Vassallo P.); bottom center) *Mesanthura* sp. (photo: Lorenti M.); bottom right) *Fistularia commersonii* at San Angelo cliff (Ischia, 14 m depth; photo: Renella G.).